


## Composition and Compositional Changes During Development: Part II



**Dr. Mark A. Ritenour**  
Indian River Research and Education Center,  
Fort Pierce

**Dr. Jeffrey K. Brecht**  
Horticultural Sciences Department, Gainesville

M.A. Ritenour

UNIVERSITY OF FLORIDA  
IFAS

1

## Major Components of Fruits and Vegetables

- **Carbohydrates**
  - the most abundant and widely distributed food component derived from plants
  - amounts vary widely
    - Leafy and stem vegetables 2 - 9%
    - Starchy roots and tubers 15 - 25
    - Citrus fruits 10 - 12
    - Dessert fruits 10 - 25

UNIVERSITY OF FLORIDA  
IFAS

2

## Major Components of Fruits and Vegetables

- **Carbohydrates**
  - The structural framework, taste and food value of a fresh commodity is related to its carbohydrate content
  - Sucrose, glucose and fructose are the main soluble (sweet) sugars in horticultural crops
  - Dessert fruits and certain vegetables, *e.g.*, sweetcorn, peas, sweetpotatoes, are relatively high in sugars

UNIVERSITY OF FLORIDA  
IFAS

3

## Major Components of Fruits and Vegetables

- **Carbohydrates**
  - Polysaccharides are the main structural components of cell walls and are important in texture and softening
    - include cellulose, hemicelluloses, and pectin
  - Starch serves as a storage carbohydrate and is organized into small grains within the cell

UNIVERSITY OF FLORIDA  
IFAS

4

## Major Components of Fruits and Vegetables

- **Carbohydrates**
  - Changes in carbohydrates after harvest of horticultural commodities are among the most important from the standpoint of quality
    - Sugar loss due to respiration
    - Conversion of starch to sugars and sugars to starch
    - Conversion of sucrose to reducing sugars
    - Solubilization and breakdown of pectin polymers to pectin fragments and galacturonic acid

UNIVERSITY OF FLORIDA  
IFAS

5



32°F      41°F      59°F

UNIVERSITY OF FLORIDA  
IFAS

6

**Major Components of Fruits and Vegetables**

- **Proteins**
  - Fruits and vegetables are relatively low compared with cereals and animal products
  - Fruits <1%
  - Leafy and stem vegetables 1-2
  - Starchy vegetables 0.5-3
  - Legumes 3-8

UNIVERSITY OF FLORIDA  
IFAS

7

**Major Components of Fruits and Vegetables**

- **Proteins**
  - Proteins are involved as enzymes catalyzing metabolic processes
  - Formation or activation of new enzymes is physiologically important in various processes
    - e.g., ripening and senescence
  - Changes in the levels and activities of enzymes due to changes in cell membrane permeability may be involved in chilling injury

UNIVERSITY OF FLORIDA  
IFAS

8

**Major Components of Fruits and Vegetables**

- **Lipids**
  - Generally low in fruits and vegetables with the exception of those commodities in which lipids serve as storage reserves
    - avocado 4-30%
    - olive 15-40%
    - tree nuts 45-65%
  - In the other horticultural crops, lipids occur mainly as components of the cell membranes, cuticle, and epidermis

UNIVERSITY OF FLORIDA  
IFAS

9

**Major Components of Fruits and Vegetables**

- **Lipids**
  - Oil content is an index of avocado maturity
  - The lipids of the cuticle and epidermis are important to the appearance of most commodities
  - The cuticle is also important in protection against water loss, pathogens and mechanical injuries
  - Lipids are involved in wound healing (suberin)

UNIVERSITY OF FLORIDA  
IFAS

10

**Major Components of Fruits and Vegetables**

- **Lipids**
  - Membrane lipids may play a role in chilling injury
    - The degree of fatty acid saturation influences membrane flexibility and may change upon exposure to chilling temperature
  - Chilling sensitive plants tend to have a high percentage of saturated fatty acids, which can undergo a phase change at chilling temperatures

UNIVERSITY OF FLORIDA  
IFAS

11

**Major Components of Fruits and Vegetables**

- **Organic acids**
  - Important in respiratory metabolism and as storage compounds
    - Organic acids are important intermediate products of metabolism. The Krebs (TCA) cycle is the main channel for the oxidation of organic acids in living cells and it provides the energy required for maintenance of cell integrity

UNIVERSITY OF FLORIDA  
IFAS

12

### Major Components of Fruits and Vegetables

- Organic acids**
  - Metabolized into amino acids, which are the building blocks of proteins
  - They can also be metabolized into many other constituents

Taiz & Zeiger, Plant Physiology, 2002

13

### Major Components of Fruits and Vegetables

- Organic acids as storage compounds**
  - Some fruits, such as lemons and limes, contain as much as 2 to 3% acid of their total fresh weight
  - Titratable acidity, specific organic acids present and their relative quantities, and other factors influencing the buffering system affect pH, which can vary from 2 to 7 among various commodities
  - Food safety implications: low pH (<4.5) inhibits microbial growth

14

### Predominant Organic Acids in Various Fruits and Vegetables

Predominant acid	Commodities
Malic	<p>Fruits: apple, apricot, banana, cherry, grape, peach, pear, plum</p> <p>Vegetables: artichoke, broccoli, carrot, cauliflower, celery cucurbits, lettuce, okra, onion</p>
Citric	<p>Fruits: lemon, orange, currant, fig, gooseberry, guava, loganberry, pineapple, pomegranate, raspberry, strawberry</p> <p>Vegetables: leafy vegetables, legumes, tomato, potato, sweetpotato</p>
Tartaric	Grape (about equal to malic)

15

### Major Components of Fruits and Vegetables

- Pigments**
  - Chlorophyll** - control of chlorophyll degradation (loss of green color) is important from a quality standpoint in both fruits and vegetables
  - Normally we wish to retard the process in vegetables and promote it in ripening fruit

16

### Major Components of Fruits and Vegetables

- Pigments**
  - Carotenoids** (yellow, orange and orange-red)
    - Very stable compounds that remain intact even when senescence is well advanced
    - Synthesis of these pigments is important during fruit development but may be masked by chlorophyll (e.g., citrus, bananas)
    - In tomato, carotenoid synthesis is concurrent with chlorophyll degradation
    - Content of B-carotene (pro-vitamin A), a major carotenoid, is important for nutrition
  - Carotenoids are important antioxidants - important for human health

17

### Major Components of Fruits and Vegetables

- Pigments**
  - Phenolics**
    - The main substrates of enzymatic browning reactions of cut or damaged tissues of apple, peach, potato, etc. upon exposure to air (O<sub>2</sub>)

18

## Major Components of Fruits and Vegetables

- **Pigments**
  - Phenolics
    - Phenolics are thought to play a role in the resistance of some immature tissues to attack by pathogens
      - pre-existing compounds
      - **phytoalexins** are formed in response to attack
    - Phenolic content is generally higher in fruits than vegetables and is higher in immature than mature fruits
      - **Astringency** in immature fruits and other tissues is related to the content of tannins
    - Phenolics are important **antioxidants**, important for human health

19

## Major Components of Fruits and Vegetables

- **Pigments**
  - **Anthocyanins** - flavonoids (red, blue and purple) are phenolic compounds
    - Water soluble, unstable glycosidic compounds that are readily hydrolyzed to free anthocyanidin or oxidized to give brown oxidation products
    - The colors of anthocyanins are influenced by vacuolar pH. Often they are confined to the cells of the epidermal layer ("blush" formed in response to sunlight)

20

## Major Components of Fruits and Vegetables

- **Volatile compounds**
  - Responsible for the characteristic **aromas** of fruits and vegetables
  - The total amount of carbon involved is much less than 1% of that evolved as CO<sub>2</sub>
  - **Ethylene** is the major volatile formed at least in climacteric crops (50-75%) yet it does not contribute to typical fruit aromas
  - Typically, **only a few key volatiles** out of 50-100 are important for the particular aroma of a given commodity

21

## Major Components of Fruits and Vegetables

- **Vitamins**
  - Fruits and vegetables are generally good sources of vitamins, which are essential in human nutrition
  - Vitamins are classified as **water-soluble** and **lipid-soluble**

22

## Water-soluble and lipid-soluble vitamins found in plants


Water-Soluble Vitamins	Lipid-soluble Vitamins
Ascorbic acid (Vit. C)	Vitamin A
Thiamin	Vitamin D
Riboflavin	Vitamin E
Niacin	Vitamin K
Vitamin B <sub>6</sub>	
Folic acid	
Vitamin B <sub>12</sub>	
Biotin	
Pantothenic acid	

23

## Major Components of Fruits and Vegetables


- **Vitamins**
  - The **water-soluble vitamins**, especially ascorbic acid (Vitamin C), are very susceptible to postharvest degradation when commodities are exposed to adverse handling and storage conditions
    - high temperature
    - low relative humidity (wilting)
    - physical damage
    - chilling injury

24




Total ascorbic acid content in strawberries stored at 1, 10 or 20°C (Nunes et al., 1998)


25




26




27



28



29



30

## Major Components of Fruits and Vegetables

### • Minerals

- **Potassium** is the most abundant mineral in fruits and vegetables (as much as 1% in parsley)
  - occurs mainly in combination with organic acids
- **Calcium** is the second most important mineral constituent
  - mainly associated with cell walls and membranes
- **Magnesium** is a component of the chlorophyll molecule
- **Phosphorus** is a constituent of proteins that are important in carbohydrate metabolism and energy transfer (i.e., ATP)
- **High nitrogen content** is often associated with reduced soluble solids content, lower acidity, and increased susceptibility to physiological disorders in fruits

31

## Conclusion

- **Composition has several important considerations in postharvest horticulture**
  - Nutritional value
  - Physiological role of constituents
  - Contributions to taste and appearance
  - Relationship to harvest and postharvest practices

32