


Principles of Postharvest Horticulture

University of Florida


1



Postharvest Deterioration and Losses

Jeff Brecht
Horticultural Sciences Department, Gainesville
Mark Ritenour
Indian River Research and Education Center, Fort Pierce


2



What is Postharvest Biology?

- A Pragmatic (**practical**) science
- Primarily deals with Perishable Commodities
- **By definition:** Postharvest = After Harvest
 - Also concerned with preharvest factors (seed source, rootstock, etc.) because they strongly influence postharvest quality. **Fruit quality is set during growth.**
 - & the harvest of the crop (e.g., when & how to harvest; maturity standards)
- Ultimately, **maximum product quality is determined (fixed) at harvest**


3



Postharvest Goals

- Harvest commodities at their **optimum maturity**
- Maintain the commodity's internal and external **quality** throughout harvest, packing, storage and distribution. **Remember, it is Alive during this process**
- Deliver the commodity to consumers at the **time and in a form (e.g., ripe, cut up, etc.) that they will purchase**


4



Historical Background

- Early postharvest practices:
 - Dried fruits, vegetables, meats, etc.
 - Fermentation of juices
 - Salting or smoking
 - Canning
- **All these methods KILL the product => product is easier to transport and store**

5




Historical Background

- Early Storage Practices:
 - **Basket** making developed by 7,000 B.C.
 - **Underground pits and silos** – 9,000 B.C. By Roman times, silos were the major means of long term storage
 - Roman era – **modified atmosphere storage** of grains
 - **Ice refrigeration** developed in 1803

6

Historical Background


- Postharvest issues became more important as cities developed (urbanization)
 - The Industrial Revolution (18th century)
 - The Problem: How to deliver (quality) fresh fruits and vegetables from areas of production to areas of consumption
- All fresh produce is alive => **A Challenge to Keep Healthy Until Consumed!**



7

Historical Background


- Early Storage Practices Continued:
 - 1819, first **Modified Atmosphere** research (France)
 - 1855 & 59, **mechanical refrigeration** invented
 - 1872, **ice refrigerated rail cars** common
 - 1889 **ammonia refrigeration** widely used for icemaking
 - 1928, **refrigerated displays** used in retail stores
 - Controlled Atmosphere (CA)** research in 1927, and in 1929, the first **commercial CA storage** (England)



8

World-Wide Perspective

- World population has reached **8.17 billion**
- Technological advances have helped us feed much of the world
 - e.g. China, India & Brazil
- Many parts of the world still in need
 - Western style, capitol intensive, methods often not suitable for an area
 - Advances have resulted in some problems (e.g., Egypt's Aswan High Dam, Calif. Central Valley)



9

World-Wide Perspective


- Food **distribution** is the main worldwide problem – not food production
 - Urbanization expected to continue at a greater rate in developing countries
 - Produce grown in these areas are the least studied
- Worldwide production (2022):
 - Cereals (durable food goods) = ~3.06 billion MT
 - Fruits, Vegetables, Melons, Roots & Tubers = ~2.76 billion MT



10

Characteristics of Perishable Commodities


- Compared to grains:
 - More subject to **deterioration after harvest** (shelf life measured in days to months vs. >1 year)
 - Relatively larger in **size** (up to 5 kg vs. < 1 g)
 - Soft textured
 - Higher water content (70 to 90+% vs. 10 to 20%)
 - Higher respiration and heat production



11

Characteristics of Fresh Produce

1. Living tissues - **High respiration and heat production**
2. **High water content** (around 90-95%) – prone to injuries and shriveling
3. Highly variable in **physiology and composition**
4. Highly subject to **pathological deterioration**



12

Postharvest Losses


- Only very rough estimates are available
 - 5 to 25% in developed countries
 - 20 to 50% in developing countries
- In the US, a large portion at the consumer level = Waste (discarded edible food)
- Loss of product = much more than just the cost of the commodity



13

Food Loss vs. Food Waste


- **Food Loss:** the food is eaten or destroyed by animals, insects or microbes; spoiled, inedible
- **Food Waste:** the food is edible but is not consumed due to poor quality or unfavorable market conditions
- *The total of loss + waste is the same (about 40%) in both developing and developed countries*



14

Other Losses


- Loss of energy & labor (e.g., during packinghouse operations)
- Loss of materials (e.g., packaging)
- Cost of reconditioning (a big reason for waste)
- Cost of waste disposal



15

Other Losses

- Loss of food value (e.g., nutrients)
- Loss of organoleptic quality
 - Color changes
 - Water loss
 - Carbohydrate changes (e.g., sugar to starch conversion)
 - Volatile changes
 - Breakdown of proteins, cell wall components, softening, etc.




16

Causes of Postharvest Loss

Internal Factors

- Respiration (metabolism)
- Compositional changes
- Morphological changes
- Physiological disorders
- General senescence




17

Causes of Postharvest Loss

Outside Factors


- Temperature
- Physical damage
- Pathogens
- Relative humidity
- Atmospheric composition
- Light
- Gravity
- Rodents and other animals
- Contamination



18

Fresh Commodities Are Still ALIVE!

- They carry out **respiration**:



19

Respiration and Shelf Life

- Respiration rate is inversely related to shelf life.

Higher respiration

=> Shorter Shelf Life



20

Respiration & Temperature

- **Temperature is the most important factor** influencing the postharvest life of a given commodity
 - Dictates the speed of chemical reactions (including respiration)
- Typically, for every 18 °F (10 °C) increase, respiration increases between 2 and 4 fold



21

Example

Temperature (°F/°C)	Shelf-Life
32/0	100
50/10	33
68/20	13
86/30	7
104/40	4



22



Affect of temperature on the quality of broccoli after just 48 h of storage at either room temperature (75°F) or in the refrigerator (40°F)



23

Compositional Changes


- Water loss
- Nutrients
- Vitamins
- Antioxidants
- Starch ⇔ Sugar
- etc.



24

Water Loss

- Besides resulting in direct loss of salable weight, it is also an **important source of quality loss**
 - Appearance quality** - wilting, shriveling, accelerated development of injuries
 - Textural quality** - loss of crispness, juiciness, etc.
 - Nutritional quality** - e.g. vitamins A & C
- Rate of water loss influenced by:
 - Environmental factors** - e.g. relative humidity
 - Anatomical factors** - stomates, hairs, etc.




25

Morphological Changes (Form & Structure)

- Because horticultural commodities are living (and sometimes still growing) they often continue development in ways that sometimes detract in quality

Changes include:


- Sprouting** (onions, tubers, root crops)
- Rooting** (onions, root crops)
- Elongation & Curvature** (asparagus, gladiolus)
- Seed Germination** (tomato, pepper, grapefruit)



26

Physiological Disorders


- Definition:**
 - “Tissue damage or breakdown not related to pathogens, insects or mechanical damage.”



27

Physiological Disorders

- Temperature**
 - High temperature injury, freezing injury, or chilling injury
- Altered atmospheric gas concentrations**
 - Reduced O₂ or elevated CO₂
- Nutrition**
 - e.g., calcium deficiency or boron toxicity




Photos courtesy of Steve Sargent.

28

Temperature is #1

- Mentioned above with respiration and physiological disorders
 - Affects all aspects of metabolism
- Temperature greatly effects water loss.
- Lower temperature also slows pathogen (human and plant) development



29

Physical Damage

- Causes the greatest amount of loss to fresh horticultural products
- Affects (among other things):
 - Respiration, ethylene production, ripening, and other metabolic processes
 - Pathogen growth and ability to invade tissue
 - Tissue discoloration






Photo courtesy of Steve Sargent

30

Pathology (decay)



- Fungi, bacteria and viruses
- Preharvest (latent) and postharvest infections
- Most postharvest infections are a result of rupturing the epidermis of the commodity

31

Relative Humidity

- Higher relative humidity slows water loss from the commodity
- High relative humidities (e.g., >95%) can weaken cartons
- Free moisture stimulates pathogen development






32

Atmospheric Composition

Modified or Controlled Atmospheres

- Altering the normal gas composition surrounding a commodity (e.g., raising or lowering O₂ or CO₂ concentrations)
- **Modified Atmospheres (MA)**
 - Passive. The commodity is placed in a gas impermeable container and the crop's respiration consumes (lowers) O₂ and gives off (increases) CO₂
- **Controlled Atmospheres (CA)**
 - Active. Gas concentrations are actively regulated using special equipment






33

Causes of Postharvest Loss

Environmental Factors



- **Light**
 - Color and morphological changes (e.g., potato greening)
- **Gravity**
 - Morphological changes (e.g., bending)
- **Rodents and other animals**
- **Contamination (food safety)**

34

Commercial Considerations



- **Preharvest factors:**
 - Cultivars & Molecular Biology
 - Nutrition & Water effects
 - Weather conditions (temperature, humidity, etc.)
 - Field sanitation (both for decay & human pathogens)

35

Commercial Considerations



- **Harvesting:**
 - Is everything ready for arrival of the harvested product?
 - Labor to harvest, grade, pack, ship, etc.
 - Materials to wash, coat, label, pack, ship, etc.
 - Best time to harvest for fresh, processing, storage?
 - Use of harvest aids

36

Commercial Considerations

- **Preliminary grading in the field:**
 - Remove unmarketable produce as soon as possible

37

Commercial Considerations

- **Packingline operations:**
 - Washing, grading, sorting, sizing, waxing, etc.
 - Each step costs \$\$\$\$ Use only if increases value of the crop







Photo courtesy of Steve Sargent

38

Commercial Considerations



- **Packaging:**
 - Protects the product, reduces water loss, orients the product, excludes light & communicates information
 - Must be economic, able to support stacking, allow ventilation (cooling), facilitate recycling or disposal at destination markets

39

Commercial Considerations



- **Postharvest development:**
 - Ethylene degreening or ripening
 - Curing

40

Commercial Considerations



- **Rapid cooling:**
 - Air cooling
 - Room
 - Forced-air
 - Hydrocooling
 - Ice Cooling
 - Top icing
 - Liquid ice injection
 - Vacuum Cooling

41

Commercial Considerations

- **Storage:**
 - Increases the cost of a product
 - Accurate temperature and RH control critical
 - CA or MA storage
- **Transportation:**
 - Water – inexpensive but slow
 - Rail – more expensive but faster
 - Truck – predominant method. Fast & reliable
 - Air – Fastest, expensive, & inconsistent scheduling and temperature control

42

Commercial Considerations

- **Marketing:**
 - Identify markets and qualities desired
 - Entire process should be geared to deliver what the consumer will buy
- **Retailing:**
 - Educate the retailer how to handle your commodity
 - Continuation of the temperature, RH and sanitation chain
 - Reconditioning?

